

lournal Collection

Reviews

Research Update

Research Tools

News & Features Books & Labware Science Jobs

reviews.bmn.com

Latest

Browse Updates Subjects Journals

Browse My Virtual Journals

Search Reviews My E-Mail Alerts

MEDLINE

Section Search

Join Login Help Feedback

Join/Login Benefits

Create virtual journals

Current Opinion in Genetics & Development

<u>History</u> Results

Current Opinion in Genetics & Development

Latest Issue **Browse Issues** Search this Journal Prices & Subscriptions

About this Journal Publishers Site

Position-effect variegation and the new biology of heterochromatin

Current Opinion in Genetics & Development

Record

[Review article] Gary H Karpen

Current Opinion in Genetics & Development 1994, 4:281-291.

<u>Text only</u>, + full figures

Publications by

Gary H Karpen

Jump to this record in Evaluated MEDLINE

Related records from Evaluated MEDLINE

Related fulltext articles on BioMedNet

Fulltext articles on BioMedNet that cite this article

Quick Site Search



BioMedNet

Advanced site search

\$

Outline

- Abstract
- Abbreviations
- Introduction
- Chromatin assembly and position-effect variegation in Drosophila
- The chromatin assembly model
- > Position-effect insulator elements and chromatin structure
- > Drosophila genes that modify position-effect variegation in trans
- Telomeric and centromeric position effects in yeast
- Nuclear organization and position-effect variegation
- Position-effect variegation and physical alterations of heterochromatic DNA
- DNA copy number reductions can be associated with position-effect variegation
- Under-replication versus somatic elimination
- Conclusions and future prospects
- Acknowledgements
- References and recommended reading
- Copyright

Abstract

(2)

Q Q

The phenomenon of position-effect variegation has long been used as evidence for the importance of chromosome position to gene expression in eukaryotes. Investigations published within the past few years demonstrate that position- effect variegation is caused by multiple mechanisms, and that direct tests of hypotheses are possible with numerous model systems.

Abbreviations

E(var) — Enhancer of variegation locus; **ORC** — origin recognition complex;

PEV -position-effect variegation;

SIR —silent information regulator;


Su(var) — Suppressor of variegation locus.

Introduction

(2)

Heitz [1] recognized that some eukaryotic genomes are divided into two cytologically distinct entities, euchromatin and heterochromatin (Fig. 1). Euchromatin contains most of the single-copy DNA and mutable genes, decondenses during interphase, and replicates throughout S phase.

Heterochromatin defies simplistic definitions, but in general it contains few mutable genes, is rich in middle-repetitive and highly-repetitive sequences (including transposons), is constitutively condensed throughout the cell cycle, and replicates late in S phase (reviewed in [2]). Position-effect variegation (PEV) was first characterized by Muller [3] as the variable, but heritable, inhibition of euchromatic gene activity when artificially juxtaposed with heterochromatin by chromosome rearrangement (<u>Fig. 1a</u>). Numerous reviews published in the past few years have summarized the interesting history of researchers' accomplishments in this field $[4][5][6][7][8][9 \bullet]$. We now recognize that position effects include a broad array of phenomena, such as heterochromatininduced inhibition of transcription and reduction in DNA copy number, telomere-induced position effects, interactions between genes on separate chromosomes (' trans -sensing effects'), and inhibition of heterochromatic genes and chromosome transmission functions (Fig. 1b). Thus, PEV provides a window for investigating the function and metabolism of heterochromatin, as well as other aspects of chromosome and nuclear organization. In this review, I will describe key developments in the field of PEV that have occurred during the past year, with special emphasis on new concepts resulting from molecular-genetic studies using tractable systems such as Drosophila and yeast. I will emphasize the viewpoint that chromatin assembly and transcriptional inactivation are not the sole mechanisms for PEV; there is mounting evidence that multiple PEV mechanisms



inversion causing PEV. A chromosome rearrangement juxtaposing euchromatin and heterochromatin causes PEV (breakpoints are indicated by open triangles). This rearrangement brings the euchromatin (shown as a line), which includes genes such as the white(w) eye-colour gene, close to centric heterochromatin causing a white $^{mottled}($ w $^m)$ or variegated phenotype. The centric heterochromatin can be separated into α (gray) and eta (diagonal bars) heterochromatin (reviewed in [75]). The lpha heterochromatin contains highly repeated satellite DNA, and some middle-repetitive elements, and is severely under- represented in dipteran polytene chromosomes. The eta heterochromatin is the 'buffer' between euchromatin and lphaheterochromatin, contains middle-repetitive transposon-like sequences and some single copy genes, and is not under- represented in polytene nuclei. Subtelomeric heterochromatin resembles $^{\sharp}$ heterochromatin in structure and can cause PEV of inserted genes [77•][79]. Eye phenotypes are shown for males with different sex chromosome constitutions (XY, XO, XYY). Dark areas indicate the normal red pigmentation (w^+ expression) and white areas indicate ommatidia that lack w ⁺activity. Removal of the predominantly heterochromatic Y chromosome (XO males) significantly reduces total genomic heterochromatin and enhances the PEV in trans(more mutant), whereas additional heterochromatin (XYY males) suppresses PEV. (b) The structure of chromosome 2 and an inversion causing PEV. Normally heterochromatic genes, such as the lightgene (lt), are moved into centromere-distal euchromatin [60]. In this case, the eye pigmentation phenotype responds to the amount of genomic heterochromatin in a 'reverse' manner to the heterochromatin-induced PEV in (a).

Return to text reference [1] [2] [3] [4] [5] [6]

Chromatin assembly and position-effect variegation in Drosophila



The chromatin assembly model



The hypothesis that chromatin compaction and transcr- iptional inactivation are the molecular mechanisms responsible for PEV arose from cytological observations on *Drosophila* polytene (endoreplicated) chromosomes (reviewed in [4]). Euchromatin juxtaposed with heterochromatin via chromosome rearrangement can display diffuse banding and high compaction normally characteristic of the heterochromatic chromocenter. This visible 'heterochromatinization' correlates with inhibition of gene function; those regions closest to the junction are most likely to appear compacted and to contain inactive genes. Elegant models (reviewed in [6]) have suggested that multimeric complexes of proteins normally present in heterochromatin are responsible for packaging large chromosomal domains in a repressed state. In these models, mass-action or self-assembly of the complexes is responsible for euchromatic 'spreading' of repressed gene activity, clonal inheritance of inactivation, and sensitivity of PEV to the dosage of heterochromatin and unlinked modifiers in the cell [10].

Although the chromatin assembly model has gained widespread acceptance, direct proof in multicellular eukaryotes, such as *Drosophila*, has been lacking (see below). In addition, new observations question the generality of previous cytogenetic characterizations of heterochromatinization. Directional spreading of chromatin assembly molecules is inconsistent with the discontinuous compaction seen upon close inspection of some variegating rearrangements [11]. Furthermore, recent studies have shown that two rearrangements present in the same cell can behave independently with respect to gene expression, compaction, and the binding of one heterochromatin-specific protein [12•][13•], discounting the hypothesis that cell-by-cell variations in gene expression are caused by differences in dosage of heterochromatinization proteins that are uniformly distributed in the nucleus. Recent findings support a role for other PEV mechanisms, such as nuclear positioning and somatic elimination, more consistent with these stochastic behaviors (see below).

Position-effect insulator elements and chromatin structure



Does chromatin structure actually change in response to PEV? Unfortunately, it has been difficult to analyze chromatin changes associated with PEV at the molecular or biochemical levels. The significance of chromatin structure in PEV has been questioned

by a comprehensive analysis of the *white* gene in the inversion $In(1)w^{m4}$. Only minor changes in DNase I sensitivity and nucleosome spacing were observed in response to this PEV rearrangement, despite extensive cytologically visible changes in polytene chromosome structure [14].

Putative chromosome domain boundary elements, such as specialized chromatin structures, can insulate chromosomally integrated genes from euchromatin-induced position effects [15] and block enhancer- mediated activation of transcription [16]. Recently, position- effect insulator elements have been identified from short DNA sequences associated with the nuclear scaffold [17], constitutive DNase I hypersensitivity $[18 \cdot \bullet]$, and transcription factor binding $[19][20 \bullet]$. Surprisingly, insulator function appears to be highly conserved during evolution. A yeast scaffold attachment region element functions as an insulator in plant cells [17], and a chicken [3] -globin constitutive

hypersensitive site insulates reporter gene function in human erythroid cells and *Drosophila* tissues [18••]. Further investigations will be required to determine whether primary sequence or secondary structure of these elements is recognized by

other species, and how much of the protein machinery is conserved.

The identification of position-effect insulator elements suggests that chromatin accessibility can influence position effects. However, a role for insulator elements in heterochromatin-induced PEV is suggested by only one study [20•], which does not address the importance of chromatin assembly directly; for example, insulator elements could act by altering the nuclear compartmentalization of a gene through attachment to the nuclear membrane or scaffold. Chromatin changes, if demonstrated to occur, could be a secondary consequence of nuclear positioning.

Drosophila genes that modify position-effect variegation in trans



If the chromatin assembly model is correct, then loci encoding trans -acting modifiers of PEV should play a role in chromatin assembly, packaging, or maintenance. As many as 120 dominant suppressors (Su(var)) or enhancers (E(var))

of variegation exist in *Drosophila melanogaster* (see [6][8] for extensive reviews). One of the best studied modifier genes is the Su(var)205 gene, which encodes the heterochromatin-binding protein HP1 [21]. The conservation of HP1 in evolutionarily distant species (mealybugs [22], *Drosophila virilis* [23], mice [24] and human [25•]), and the lethality of HP1 null alleles [26•], has suggested that HP1-like proteins are important for cell viability and/or development. However, the recessive lethal phenotypes of HP1 null animals are consistent with diverse roles for the protein, including activation of essential heterochromatic genes [27], chromosome transmission, or regulation of transposable elements [26•].

HP1 shares the 52 amino acid 'chromo domain' [28] or 'HP1/Pc box' [23] with the Polycomb protein (Pc), which regulates important developmental events by repressing euchromatic homeotic gene expression. A link between PEV and homeotic gene repression has been substantiated by the exciting demonstration that regulatory regions that respond to Pc group repression can induce variegation of an adjacent *white* gene [29••]. The fact that neither HP1 nor Pc proteins themselves bind DNA [30••], and the dose-dependent phenotypes associated with mutations at each locus [21][28], have led to the speculation that both proteins are involved in the assembly of different multimeric complexes that maintain repressed gene activity during development (reviewed in [31•]).

Does the chromo domain function to compact chromatin? Molecular dissection of HP1 protein functions indicates that nuclear localization and heterochromatin association functions map outside the chromo domain, but this study does not directly address

the function of the chromo domain [32]. Monitoring the binding of mutant Pc proteins to euchromatic polytene chromosome sites has elegantly demonstrated that the chromo domain is necessary for proper Pc distribution [30••]. These results support the hypothesis that the chromo domain promotes protein–protein interactions, presumably between Pc, or HP1, and DNA-binding proteins that are responsible for localization of the complexes to specific chromosomal sites. As HP1 and Pc display distinctly different chromosomal distributions, their binding specificity must be encoded by minor differences between the chromo domains, or by another part of the amino termini. It is still unclear whether these complexes accomplish gene repression by higher-order chromatin compaction, nuclear compartmentalization, or other mechanisms (see below) [31•].

The involvement of proteins such as HP1 in PEV is likely to tell only part of the story. Only a few of the products from the hundreds of modifier loci have been analyzed by molecular cloning, and they vary in structure and potential function.

Cloned genes encoding modifiers include Su(var)205 (HP1), Su(var)(3)7 (function unknown, the protein contains unusually spaced zinc fingers [33]), modulo (a DNA-binding protein [34•]), and Su(var)(3)6 (the protein phosphatase 1 catalytic subunit [35]). The identification of recessive PEV modifier mutations, whose phenotypes are not dose dependent [36], opens up the possibility that this new class of genes can be identified by direct genetic screening. The biological functions of even those genes that have been cloned are still obscure. A number of molecular mechanisms could be carried out by proteins associated with DNA, including chromatin compaction, but also nuclear localization, replication, nuclease activity, transposition and recombination. Our understanding of the diversity of mechanisms acting on heterochromatin will be greatly enhanced by direct demonstrations of the biological functions of the proteins. The isolation of readily clonable Pelement induced Su(var) and E(var) alleles $[37 \bullet 1][38]$ should greatly facilitate this important undertaking.

Telomeric and centromeric position effects in yeast



The relevance of <code>Drosophila</code> PEV models to other organisms has been validated by recent studies in yeasts (reviewed in $[9 \bullet][39 \bullet \bullet][40 \bullet]$). In <code>Saccharomyces</code> <code>cerevisiae</code>, inactive mating-type loci (<code>HML</code> and <code>HMR</code>) and genes inserted near telomeres display heritable, but reversible, repression reminiscent of the PEV phenotypes seen in multicellular eukaryotes <code>[41]</code>. As for PEV in <code>Drosophila</code>, telomeric silencing in yeast is directional and the extent of the silenced region can be modified by altering the dosage of the product of an unlinked locus (<code>SIR3</code>, a silent information regulator) <code>[42 \bullet]</code>. However, the yeast position effects include $\sim 2-3$ orders of magnitude less <code>DNA</code>, and inactivation of telomeric genes is normally less frequent, than heterochromatin-induced PEV in <code>Drosophila</code>.

The most direct biochemical and molecular evidence that PEV involves chromatin structure changes comes from detailed studies in S. cerevisiae and Schizosaccharomyces pombe. Resistance to methyltransferase activity [43][44], unusual nucleosome structures [45], suppression of silencing by histone H3 and H4 mutations [46], and hypoacetylation of histones [47] at telomeres and HM loci show that detectable chromatin changes are correlated with the silenced phenotype. S. pombe centromeric regions have been shown recently to induce PEV on inserted genes [48•], and the centromere central core is associated with unusual nucleosome spacing [48•][49].

The primary cause of chromatin changes and PEV in yeast is still unknown, but progress in this area is rapid. The mapping of an origin of replication and a sequence required for silencing to the same 138 bp HMR element has forged a link between replication and silencing $[50 \bullet]$. The origin recognition complex (ORC) binds autonomously replicating sequences and is required for $in \ vivo$ origin function. Elegant genetic $[51 \bullet \bullet][52 \bullet \bullet]$ and biochemical $[53 \bullet \bullet]$ analyses have demonstrated that the ORC is essential for silencing. ORC binding alone, rather than the initiation of replication, appears to be required to recruit proteins encoded by the SIR s to HM loci $[54 \bullet]$. It is still unclear whether ORC and subsequent SIR binding are required for establishment and/or maintenance of the repressed state $[51 \bullet \bullet][52 \bullet \bullet][53 \bullet \bullet]$. However, recent experiments investigating the cell-cycle dependence of activation of a telomere-silenced gene have suggested that replication may be necessary to reverse silencing (OM Aparicio, DE Gottschling, personal communication). The role of the RAP1 protein in PEV and localizing telomeres to the nuclear membrane will be discussed below.

Nuclear organization and position-effect variegation



Interphase nuclei display a characteristic organization (the Rabl orientation [55]). In general, centromeres and telomeres are clustered and are found associated with opposite poles of the nuclear envelope [56], their positioning and clustering being regulated during the cell cycle [57•][58•]. A number of studies published in the past few years have suggested that the positions of genes within the nucleus, not just within the euchromatic or heterochromatic regions of chromosomes, are important for normal expression (reviewed in [59•]). This constitutes a major change in our perception of the mechanisms responsible for PEV; genetic systems for investigating the functional significance of nuclear positioning are emerging rapidly.

It is difficult to explain the effects of distant structural changes on some PEV systems, and the discontinuous compaction of euchromatin associated with some rearrangements (see [12•] and text above), as the result of spreading chromatin compaction or decompaction along the chromosome. Tantalizing examples of unusual PEVs in *Drosophila* have suggested that heterochromatic compartment(s) exist within the nucleus and are important for normal gene function. The rare genes present in Drosophila heterochromatin display a 'reverse' position effect, that is, their function is inhibited when moved into euchromatin by chromosome rearrangement (Fig. 1b). PEV of heterochromatic genes, however, appears to be caused by increased distance from major heterochromatic blocks, rather than simple juxtaposition with euchromatin [60][61]. Interestingly, modifiers of PEV (such as changing the dosage of total genomic heterochromatin, or trans -acting genes) in general have opposite effects on PEV of euchromatic genes and PEV of heterochromatic genes [27] and chromosome transmission [62•] (Fig. 1b). Recent genetic studies have suggested that PEV of euchromatic genes is also influenced by nuclear position. Autosomal rearrangements that revert brown Dominant PEV move this locus (the normally euchromatic brown gene plus the adjacent insertion of a large block of heterochromatin) to the autosomal tips [63•]. Finally, PEV is enhanced by the removal of terminal sequences from a Drosophila minichromosome, even for terminal deficiency breakpoints up to 100 kb from the affected euchromatic gene [64][65].

These examples are consistent with models involving 'looping' of affected regions into heterochromatic (centromeric and/or telomeric) nuclear domains formed by Rabl configuration clustering. Looping into the heterochromatic domain would repress euchromatic gene function [63•], but would be essential for the function of heterochromatic genes (Fig. 2). Further investigations are required to test this hypothesis, such as cytological examination of the positions of ectopic heterochromatic genes in interphase nuclei [66] under variegating and non-variegating circumstances. It is worth noting that transformation experiments involving ribosomal genes demonstrate that their normal heterochromatic location is not required for RNA polymerase I transcription, nucleolus formation, or X-Y meiotic pairing [67][68], suggesting that these processes are not sensitive to either nuclear or chromosomal position.




Fig. 2.A nuclear positioning model for PEV. (a)An interphase nucleus with heterochromatic domains formed by telomeric and centromeric clustering (Rabl configuration, see text). A representative chromosome is shown, with a heterochromatic gene indicated by a black box and a euchromatic gene indicated by an open box (see Fig. 1 for other symbols). (b,c)A rearrangement of the chromosome is shown, with the resulting gene expression being determined by the position of the genes within the nucleus. (b) Looping into the euchromatic domain would be essential for the expression of the euchromatic gene, but would repress the expression of the heterochromatic gene [60][61]. (c) Looping into the heterochromatic domain would be essential for the expression of the heterochromatic gene, but would repress euchromatic gene expression [63•]. Once located to the heterochromatic compartment, euchromatic genes could be inactivated by exclusion of transcriptional or replication machinery from the compartment, domain-induced changes in chromatin structure, or other types of modification (see text). In this model, the probability that a particular site would loop into a heterochromatic compartment would depend on the amount of heterochromatin at the site, the total amount of heterochromatin in the cell, and the distance between the site and the compartment.

Return to text reference [1]

Results from studies of yeast position effects have suggested that telomere associations with the nuclear membrane are an important component of silencing and may be responsible for observed changes in chromatin structure. RAP1 protein binds to telomeric repeats [69], is responsible for telomere-telomere and telomere- -nuclear membrane associations (reviewed in [59•]) and is required for telomeric silencing [70][71•]. Furthermore, the SIR3 and SIR4 products are localized to the nuclear periphery, and mutations in either gene result in both loss of telomere-membrane associations and derepression of telomeric and HM silencing [72•]. The recent discovery that RAP1 dependent and SIR2,3,4 -dependent silencing can be induced by C $_{1-3}$ A terminal repeats inserted up to 400 kb from the telomere argues for a critical role for the repeats, rather than telomere location per se (JB Stavenhhagen, VA Zakian, personal communication). The possibility that internal repeats accomplish silencing by looping to the telomere/nuclear membrane compartment has been suggested by the RAP1 -dependence and the observation that C $_{\mathrm{1-3}}$ A silencing is stronger at telomere-proximal locations. PEV associated with genes inserted at the S. pombe centromere requires a centromeric location, and not just central core sequences [48.]. Fission yeast centromeres are located at the nuclear periphery in interphase [49], but a functional link between the centromeric position effect and nuclear position has not been demonstrated. Perhaps the primary event in yeast silencing is localization to the nuclear membrane (dependent on SIR3, 4 / RAP1?), followed by directional propagation of (SIR1 , 2 ,/ histone H4 / ARD1 / NAT1 -mediated ?) chromatin changes. Proof that telomere, HM or centromere associations with the membrane are sufficient for initiation will require the demonstration that returning genes to the periphery in rap1, sir3 or sir4 mutants restores silencing.

Position-effect variegation and physical alterations of heterochromatic DNA

(2)

DNA copy number reductions can be associated with positioneffect variegation **(2)**

The genetic and cytological characteristics of PEV are consistent with molecular mechanisms other than transcriptional inactivation via chromatin compaction and/or alterations in nuclear positioning. Physical modifications to the DNA, such as reduced gene copy number, could play a role in PEV, and provide an attractive explanation for the heritability of the repressed state through many cell divisions.

In dipteran polytene chromosomes, heterochromatic DNA copy number is reduced (underrepresented) 64–1000-fold, with respect to the euchromatin [73]. Studies of the copy number of sequences in the *Drosophila* minichromosome Dp1187 ($\underline{Fig. 3}$) in polytene cells has provided strong support for the inclusion of adjacent euchromatin in heterochromatin-associated underrepresentation [74]. Although changes in euchromatic copy number were sufficient to account for phenotypic ($yellow^+$) variegation ($\underline{Fig. 3}$), an additional effect on transcription could not be ruled out [74].




Fig. 3.Structure and under-representation of the *Drosophila*minichromosome *Dp1187*. **(a)**The molecular structure of the 1300 kb minichromosome *Dp1187*. The thin line represents euchromatin, which contains genes normally present on the tip of the X, including the *yellow* +body-colour gene located at - 20 kb. 0 kb marks the euchromatin-

of the X, including the *yellow* *body-colour gene located at - 20 kb. 0 kb marks the euchromatin-heterochromatin junction. The lightly tinted box represents the centric (α) heterochromatin, and the black box indicates the position of an 'island of complex DNA', which is a block of middle-repetitive and/or single copy sequences that is found between satellite-containing regions [74][77•]. Two other islands are present in the *Dp1187*heterochromatin (GH Karpen, unpublished data). **(b)**The DNA copy number for different regions of the minichromosome, relative to the same region on the normal X (% representation), is shown for XO salivary gland polytene nuclei [74][77•].

Return to text reference [1] [2] [3] [4] [5] [6] [7]

The fact that some PEV rearrangements display euchromatic under-representation $\underline{[74][76]}$, whereas others do not $\underline{[14][77^\bullet]}$, suggests that heterochromatin is functionally diverse and that multiple mechanisms are responsible for PEV. The under-representation of juxtaposed euchromatin probably depends on the usual representation of heterochromatic regions present at the junction. For example, rearrangements involving β - heterochromatin (Fig. 1; reviewed in [75]) may not involve euchromatic under-representation, because at least some β - heterochromatin is known to be fully represented in polytene nuclei [78]. A striking example of the diversity of PEV mechanisms comes from a comparison of two different regions of the <code>Dp1187</code> minichromosome.

The strong inhibition of rosy ⁺ gene expression displayed by P-element constructs inserted in the subtelomeric heterochromatin (<u>Fig. 3</u>) involves only minor under-representation and is most probably caused by transcriptional inactivation [<u>77•</u>]. The subtelomeric insertions are located in a region rich in middle-repetitive elements, which is only 220 kb from the extensively under-represented yellow ⁺ gene adjacent to centric heterochromatin. Determining the roles of nuclear position and chromatin compaction in this and other telomeric position effects in *Drosophila* (see [<u>77•][79]</u> for references) requires further investigation.

Under-replication versus somatic elimination



> What is the molecular mechanism responsible for heterochromatin- associated changes in DNA copy number? In S. cerevisiae, early replication origins become utilized late in S phase when inserted near telomeres [80•]. Under-representation in Drosophila could arise during polytenization because replication of heterochromatin is inhibited [81] or delayed [82•] (the 'under-replication' model), for example by packaging into a repressed chromatin structure, or inclusion in a nuclear compartment incompatible with normal replication. Alternatively, heterochromatic DNA may be physically removed during polytenization ('elimination'), and perhaps even in diploid somatic cells. DNA elimination has been documented in distantly related eukaryotic species (ciliates, nematodes, crustaceans, dipterans, and vertebrates), and frequently involves heterochromatin and polytenization (reviewed in [83][84.1[85]).

> Recent experiments utilizing the Dp1187 minichromosome have suggested that elimination, rather than under-replication, is responsible for heterochromatic under-representation. DNA from a region of *Dp1187* displaying a gradient of under-representation in salivary gland nuclei (near 0 kb in Fig. 3) was analyzed by two- dimensional electrophoresis [86•]. Stalled replication forks, predicted by the under-replication model to be present in 33% of the molecules, were not found. Positive evidence for qualitative changes in chromosome structure associated with heterochromatin comes from recent pulsed-field analyses. Severely shortened chromosomes that lack the satellite DNAs normally present in Dp1187 (Fig. 3) appear in polytenized tissues and are likely to retain only Dp1187 euchromatin ([87•]; GH Karpen, AC Spradling, unpublished data).

> Transposon-like elements (e.g. Tecs) are removed during macronuclear polytenization in Euplotes and Oxytricha [84••][88]. Perhaps transposon-like sequences in the 'islands of complex DNA' (Fig. 3) are responsible for under-representation in Drosophila . However, the elucidation of the exact mechanism of under-representation requires further experimentation, including in-depth structural analysis of developmentally altered molecules. The detailed restriction map of Dp1187 heterochromatin (Fig. 3) and the isolation of a large number of minichromosome deletion derivatives (GH Karpen, unpublished data) will help in the testing of models by mapping cis sequences responsible for PEV and under-representation. Although exact mechanisms are not currently understood, the functional ramifications of heterochromatin undergoing developmentally regulated covalent changes in multicellular eukaryotes are numerous [4][85][87•], and warrant further exploration.

Conclusions and future prospects



The past year has seen renewed interest in PEV, yielding new information about PEV mechanisms and components, and producing manipulatable systems that hold promise for increasing our understanding in the immediate future. Major advances have come from detailed molecular-genetic analyses of telomeric and centromeric position effects and mating-type locus silencing in the yeasts S. cerevisiae and S. pombe . Further investigations should begin to elucidate the complex interaction between DNA replication and the establishment or maintenance of the silenced state, reveal chromatin structure and protein-component changes associated with position effects, and yield insight into the role of telomere associations with the nuclear envelope.

However, heterochromatin in multicellular eukaryotes, such as Drosophila and humans, is structurally and functionally more complex than that found in yeasts. It is encouraging that 'simplified' systems have been developed in recent years that allow heterochromatin and PEV to be studied with the specificity and directness necessary for progress. For example, correlating functions with the molecular structure of heterochromatin (e.g. Dp1187) will help in the elucidation of the nature of the cis sequences responsible for inducing PEV and the responsiveness of the variegating domains. From studies of brown and light variegation, and readily clonable modifier loci, we look forward to direct analyses of chromatin changes in affected tissues, molecular-cytological proof of the importance of nuclear positioning, and extensive analysis of the molecular and biochemical functions of modifier loci.

It is important to emphasize that direct proof of a primary role for chromatin assembly, the most widely accepted model for heterochromatin-induced PEV, is lacking in multicellular eukaryotes. Recent data, reviewed here, suggest that multiple mechanisms are responsible for the group of phenomena we call PEV, reflecting the structural and functional diversity of heterochromatin. It is likely that a number of the Su(var) and E(var) loci encode products that directly (e.g. DNA-binding proteins or packaging components, such as histones) or indirectly (e.g. regulatory phosphatases and kinases) influence chromatin structure. Investigators need to be more cautious, however, in assigning chromatin assembly or maintenance functions ad hoc to genes whose mutant products modify the phenotypes associated with PEV rearrangements. We cannot be blind to the exciting possibility that modifiers of PEV may function to regulate heterochromatic DNA elimination or inheritance functions, encode boundary functions that separate higher-order chromosome domains, regulate the position of domains within the nucleus, or control (as yet) undiscovered mechanisms responsible for the behavior and function of the mysterious entity known as heterochromatin.

Acknowledgements



I thank Dan Gottschling, Mike McKeown, Allan Spradling, and Barbara Wakimoto for critical comments and discussions. Helpful corrections and comments from Kevin Cook, Joan Esnayra, Huong Le, Terence Murphy and Janice Wahlstrom are gratefully acknowledged. Work in the author's laboratory is supported by NIH grant R01 HG00747, a Pew Scholars Fellowship, and Cancer Center Support Grant CA14195.

References and recommended reading





> Papers of particular interest, published within the annual period of review, have been highlighted as: of special interest.

of outstanding interest.

1. Heitz E:

Das Heterochromatin der Moose.

I Jahrb Wiss Botanik 1928, 69: 762-818.

Return to citation reference [1]

2. John B: The Biology of Heterochromatin. In Heterochromatin: Molecular and Structural Aspects Edited by Verma RS. Cambridge: Cambridge University Press, 1988, 1-147.

Return to citation reference [1]

3. Muller HJ:

Types of Visible Variations Induced by X-Rays in Drosophila.

J Genet 1930, 22: 299-334.

Return to citation reference [1]

4. Spradling AC, Karpen GH:

Sixty Years of Mystery.

Genetics 1990, 126: 779-784. MEDLINE Cited by

Return to citation reference [1] [2] [3]

5. Eissenberg JC, Elgin SC:

Boundary Functions in the Control of Gene Expression.

Trends Genet 1991, 7: 335-340. ScienceDirect MEDLINE Cited by

Return to citation reference [1]

6. Grigliatti T:

Position-Effect Variegation — an Assay for Nonhistone Chromosomal Proteins and **Chromatin Assembly and Modifying Factors.**

Methods Cell Biol 1991, 35: 587-627. MEDLINE Cited by

Return to citation reference [1] [2] [3]

7. Henikoff S:

Position Effect and Related Phenomena.

Curr Opin Genet Dev 1992, 2: 907-912. MEDLINE Cited by

Return to citation reference [1] [2]

8. Reuter G, Spierer P:

Position Effect Variegation and Chromatin Proteins.

Bioessays 1992, 14: 605-612. MEDLINE Cited by

Return to citation reference [1] [2]

9. • Rivier DH, Rine J:

Silencing: the Establishment and Inheritance of Stable, Repressed Transcription

Curr Opin Genet Dev 1992, 2: 286-292. MEDLINE Cited by

An excellent review of telomeric and mating-type loci silencing in Saccharomyces cerevisiae , emphasizing the heritability of altered chromatin structure at these regions, and the role of replication in establishing the repressed state.

Return to citation reference [1] [2]

10. Locke J, Kotarski MA, Tartof KD:

Dosage-Dependent Modifiers of Position Effect Variegation in Drosophila and a Mass Action Model that Explains their Effect.

Genetics 1988, 120: 181- 198. MEDLINE Cited by

Return to citation reference [1]

11. Belyaeva ES, Zhimulev IF:

Cytogenetic and Molecular Aspects of Position Effect Variegation in Drosophila. III. Continuous and Discontinuous Compaction of Chromosomal Material As a Result of Position Effect Variegation.

Chromosoma 1991, 100: 453-466. MEDLINE Cited by

Return to citation reference [1]

12. • Belyaeva ES, Demakova OV, Umbetova LF, Zhimulev IF:

Cytogenetic and Molecular Aspects of Position Effect Variegation in *Drosophila*. V. Heterochromatin-Associated Protein HP1 Appears in Euchromatic Chromosomal Regions that Are Inactivated as a Result of Position-Effect Variegation.

Chromosoma 1993, 102: 583-590. MEDLINE Cited by

This study demonstrates that compacted regions of polytene chromosomes contain HP1 protein; however, two PEV rearrangements can display different HP1-binding behaviors in the same cell. This result suggests that PEV can be associated with stochastic behaviors of individual chromosomes, rather than the protein content of individual cells.

Return to citation reference [1] [2] [3]

13. • Bishop CP:

Evidence for Intrinsic Differences in the Formation of Chromatin Domains in *Drosophila* melanogaster.

Genetics 1992, 132: 1063-1069. MEDLINE Cited by

Analysis of the PEV phenotypes of two rearrangements in the same cell demonstrates independent responses, again arguing (see $[12 \cdot]$) against the hypothesis that cell-by-cell variegation arises because cells differ in the dosage of 'heterochromatization' proteins.

Return to citation reference [1]

14. Hayashi S, Ruddell A, Sinclair D, Grigliatti T:

Chromosomal Structure Is Altered by Mutations that Suppress or Enhance Position Effect Variegation.

Chromosoma 1990, **99**: 391– 400. MEDLINE Cited by

Return to citation reference [1] [2]

15. Kellum R, Schedl P:

A Position-Effect Assay for Boundaries of Higher Order Chromosomal Domains. *Cell* 1991, **64**: 941–950. MEDLINE Cited by

Return to citation reference [1]

16. Kellum RB, Schedl P:

A Group of scs Elements Function As Domain Boundaries in an Enhancer-Blocking Assay.

Mol Cell Biol 1992, 12: 2424-2431. MEDLINE Cited by

Return to citation reference [1]

17. Allen GC, Hall JGE, Childs LC, Weissinger AK, Spiker S, Thompson WF:

Scaffold Attachment Regions Increase Reporter Gene Expression in Stably Transformed Plant Cells.

Plant Cell 1993, 5: 603-613. MEDLINE Cited by

Return to citation reference [1] [2]

18. •• Chung JH, Whiteley M, Felsenfeld G:

A 5' Element of the Chicken Beta-Globin Domain Serves As an Insulator in Human Erythroid Cells and Protects Against Position Effect in *Drosophila*.

Cell 1993, 74: 505-514. MEDLINE Cited by

A striking demonstration of the evolutionary conservation of elements that can insulate genes from PEV. A sequence 5' to the chicken beta-globin gene was shown to protect genes from positive or negative influences of adjacent sequences in both a human erythroid cell line and Drosophila germ-line transformants. In the human cell line, insulation involved alteration of the promoter's chromatin structure.

Return to citation reference [1] [2]

19. Geyer PK, Corces VG:

DNA Position-Specific Repression of Transcription by a *Drosophila*Zinc Finger Protein. Genes Dev 1992, 6: 1865–1873. MEDLINE Cited by

Return to citation reference [1]

20. • Roseman RR, Pirrotta V, Geyer PK:

The Su(Hw) Protein Insulates Expression of the *Drosophila melanogasterwhite*Gene from Chromosomal Position-Effects.

EMBO J 1993, 12: 435-442. MEDLINE Cited by

This paper reports that a transcription factor (Su(Hw)) binding site can protect a reporter gene from position effects, though heterochromatin-induced PEV was only partially suppressed. The Su(Hw) locus acted as a dominant enhancer of position effects displayed

by these constructs, that is less Su(Hw) protein resulted in less reporter gene expression. The authors suggest that the repressive behavior of this binding site could explain the mutagenic effects of transposon insertions, especially in genes with large regulatory regions.

Return to citation reference [1] [2]

21. Eissenberg JC, Morris GD, Reuter G, Hartnett T:

The Heterochromatin-Associated Protein HP-1 Is an Essential Protein in *Drosophila* with Dosage-Dependent Effects on Position-Effect Variegation.

Genetics 1992, 131: 345-352. MEDLINE Cited by

Return to citation reference [1] [2]

22. Epstein H, James TC, Singh PB:

Cloning and Expression of *Drosophila*HP1 Homologs from a Mealybug, *Planococcus citri*. *J Cell Sci* 1992, **101**: 463-474. <u>Cited by</u>

Return to citation reference [1] [2]

23. Clark RF, Elgin SC:

Heterochromatin Protein 1, a Known Suppressor of Position-Effect Variegation, Is Highly Conserved in *Drosophila*.

Nucleic Acids Res 1992, 20: 6067-6074. MEDLINE Cited by

Return to citation reference [1] [2] [3]

24. Hamvas RM, Reik W, Gaunt SJ, Brown SD, Singh PB:

Mapping of a Mouse Homolog of a Heterochromatin Protein Gene to the X Chromosome.

Mamm Genome 1992, 2: 72-75. MEDLINE Cited by

Return to citation reference [1] [2]

25. • Saunders WS, Chue C, Goebl M, Craig C, Clark RF, Powers JA, Eissenberg JC, Elgin SC, Rothfield NF, Earnshaw WC:

Molecular Cloning of a Human Homologue of *Drosophila*Heterochromatin Protein HP1 Using Anti-Centromere Autoantibodies with Anti-Chromo Specificity.

J Cell Sci 1993, 104: 573-582. Cited by

This most recent demonstration of the conservation of HP1 proteins in widely divergent species (see also [22][23][24]) is suggestive of an important role for this heterochromatin-binding protein. A number of other proteins of similar molecular weight react to the 'anti-chromo' antibody, suggesting that a protein family exists in humans. The functional significance of the structural homology with HP1 awaits further analyses, including a description of the chromosomal distribution in humans.

Return to citation reference [1]

26. • Eissenberg JC, Hartnett T:

A Heat Shock-Activated cDNA Rescues the Recessive Lethality of Mutations in the Heterochromatin-Associated Protein HP1 of *Drosophila melanogaster*.

Mol Gen Genet 1993, **240**: 333-338. MEDLINE Cited by

HP1 fusion constructs were demonstrated to rescue the recessive lethality associated with Su(var)205 null mutations, confirming that HP1 is essential for viability. A possible role for HP1 in chromosome transmission is suggested because of the observed aneuploidy in mutant animals, but responsibility for any of a number of other functions could not be excluded.

Return to citation reference [1] [2]

27. Hearn MG, Hedrick A, Grigliatti TA, Wakimoto BT:

The Effect of Modifiers of Position-Effect Variegation on the Variegation of Heterochromatic Genes of *Drosophila melanogaster*.

Genetics 1991, 128: 785-797. MEDLINE Cited by

Return to citation reference [1] [2]

28. Paro R. Hogness DS:

The Polycomb Protein Shares a Homologous Domain with a Heterochromatin-Associated Protein of *Drosophila*.

Proc Natl Acad Sci U S A 1991, 88: 263-267. MEDLINE Cited by

Return to citation reference [1] [2]

29. •• Fauvarque MO, Dura JM:

polyhomeoticRegulatory Sequences Induce Developmental Regulator-Dependent Variegation and Targeted P-Element Insertions in Drosophila.

Genes Dev 1993, 7: 1508-1520. MEDLINE Cited by

This paper describes a new type of position effect, termed developmental regulator effect variegation (DREV). Reporter gene expression was dramatically repressed by adjacent polyhomeotic (ph a Polycomb group member) regulatory sequences. The variegation was shown to be sensitive to the cellular dosage of Pc and ph. However, direct relevance to heterochromatin-induced PEV is questioned by the insensitivity of DREV to PEV modifiers. The targeting of polyhomeotic P element insertions to Pc group protein binding sites is described, and the authors discuss the possibility that multimeric protein complexes are responsible for this behavior.

Return to citation reference [1]

30. •• Messmer S, Franke A, Paro R:

Analysis of the Functional Role of the Polycomb Chromo Domain in Drosophila melanogaster.

Genes Dev 1992, 6: 1241-1254. MEDLINE Cited by

In this elegant study, the 'chromo domain' is demonstrated to play a role in the chromosomal distribution of the Polycomb protein. The lack of DNA-binding capability for Pc suggests that chromosomal localization is mediated by chromo domain dependent protein-protein interactions with at least one DNA-binding protein. Given the different distributions of HP1 and Pc, the specificity of binding is considered to be encoded by either minor differences in the composition of the chromo domains, or by other parts of the proteins.

Return to citation reference [1] [2]

31. • Paro R:

Mechanisms of Heritable Gene Repression During Development of Drosophila.

Curr Opin Cell Biol 1993, **5**: 999–1005. MEDLINE Cited by

A well written, detailed review of Polycomb group genes and how they act to repress inappropriate gene activity and promote stable inheritance of the determined state. The author discusses two molecular models for Pc group complex action, the older chromatin compaction model, and a new model formally similar to nuclear compartmentalization, in which the multimeric complexes simply exclude transcriptional activators from a chromosomal region.

Return to citation reference [1] [2]

32. Powers JA, Eissenberg JC:

Overlapping Domains of the Heterochromatin-Associated Protein HP1 Mediate Nuclear Localization and Heterochromatin Binding.

J Cell Biol 1993, 120: 291-299. MEDLINE Cited by

Return to citation reference [1]

33. Reuter G, Giarre M, Farah J, Gausz J, Spierer A, Spierer P:

Dependence of Position-Effect Variegation in Drosophilaon Dose of a Gene Encoding an **Unusual Zinc-Finger Protein.**

Nature 1990, 344: 219-223. MEDLINE Cited by

Return to citation reference [1]

34. • Garzino V, Pereira A, Laurenti P, Graba Y, Levis RW, Le PY, Pradel J:

Cell Lineage-Specific Expression of modulo, a Dose- Dependent Modifier of Variegation in Drosophila.

EMBO J 1992, 11: 4471-4779. MEDLINE Cited by

The first identification of a bona fide DNA-binding protein that is a dominant suppressor of PEV in Drosophila . The developmental time of appearance and nuclear distribution of the modulo protein, and genetic interactions between modulo mutations and Su(var)205, suggest that modulo may be part of a multimeric complex that includes HP1. However, modulo protein is present at greater than 100 euchromatic sites, in addition to the colocalization with HP1 to the polytene chromocenter.

Return to citation reference [1]

35. Dombradi V, Cohen PT:

Protein Phosphorylation Is Involved in the Regulation of Chromatin Condensation **During Interphase.**

FEBS Lett 1992, 312: 21-26. ScienceDirect MEDLINE Cited by

Return to citation reference [1]

36. Sinclair DA, Ruddell AA, Brock JK, Clegg NJ, Lloyd VK, Grigliatti TA:

A Cytogenetic and Genetic Characterization of a Group of Closely Linked Second Chromosome Mutations that Suppress Position-Effect Variegation in Drosophila melanogaster.

Genetics 1992, 130: 333-344. MEDLINE Cited by

Return to citation reference [1]

37. • Dorn R, Szidonya J, Korge G, Sehnert M, Taubert H, Archoukieh E, Tschiersch B, Morawietz H, Wustmann G, Hoffmann G, Reuter G:

P Transposon-Induced Dominant Enhancer Mutations of Position-Effect Variegation in Drosophila melanogaster.

Genetics 1993, 133: 279-290. MEDLINE Cited by

A large number of E(var) and Su(var) mutations were isolated after P element mediated mutagenesis. In contrast to earlier (chemical and irradiation) mutagenesis experiments, P element mediated mutagenesis yielded many more E(var) s than Su(var) s. This collection of mutations will facilitate the molecular analysis of significantly more modifiers of PEV.

Return to citation reference [1]

38. Clegg NJ, Whitehead IP, Brock JK, Sinclair DA, Mottus R, Stromotich G, Harrington MJ,

A Cytogenetic Analysis of Chromosomal Region 31 of Drosophila melanogaster. Genetics 1993, 134: 221-230. MEDLINE Cited by

Return to citation reference [1]

39. •• Laurenson P, Rine J:

Silencers, Silencing, and Heritable Transcriptional States.

Microbiol Rev 1992, 56: 543-560. MEDLINE Cited by

A scholarly, thorough review of silencing systems and mechanisms emphasizing the history and current understanding of silent mating loci in yeast. Interesting comparisons are made with heritable silencing in other organisms.

Return to citation reference [1]

40. • Sandell LL, Zakian VA:

Telomeric Position Effect in Yeast.

Trends Cell Biol 1992, 2: 10–14. ScienceDirect Cited by
A concise review describing the cis and trans components of telomeric PEV in yeast. The review discusses and contrasts three models: unusual chromatin structure, occlusion of transcription factors, and nuclear positioning.

Return to citation reference [1]

41. Gottschling DE, Aparicio OM, Billington BL, Zakian VA:

Position Effect at S. cerevisiaeTelomeres: Reversible Repression of Pol II Transcription.

Cell 1990, 63: 751-762. MEDLINE Cited by

Return to citation reference [1]

42. • Renauld H, Aparicio OM, Billington BL, Chhablani SK, Gottschling DE:

Silent Domains Are Assembled Continuously from the Telomere and Are Defined by Promoter Distance and Strength, and by SIR3 Dosage.

Genes Dev 1993, 7: 1133-1145. MEDLINE Cited by

The amount of silenced DNA at yeast telomeres was shown to be affected by the strength of URA3 basal expression and the dosage of the SIR3 protein. Directional repression of adjacent genes suggested telomere to centromere propagation of the silenced domain. An interesting model is proposed in which telomeric location of the key regulator SIR3 could create negative feedback regulation of telomeric silencing.

Return to citation reference [1]

43. Gottschling DE:

Telomere-Proximal DNA in Saccharomyces cerevisiaeIs Refractory to Methyltransferase Activity in Vivo.

Proc Natl Acad Sci U S A 1992, 89: 4062-4065. MEDLINE Cited by

Return to citation reference [1]

44. Singh J. Klar AJ:

Active Genes in Budding Yeast Display Enhanced in VivoAccessibility to Foreign DNA Methylases: a Novel in VivoProbe for Chromatin Structure of Yeast.

Genes Dev 1992, 6: 186-196. MEDLINE Cited by

Return to citation reference [1]

45. Wright JH, Gottschling DE, Zakian VA:

SaccharomycesTelomeres Assume a Non-Nucleosomal Chromatin Structure.

Genes Dev 1992, 6: 197-210. MEDLINE Cited by

Return to citation reference [1]

46. Aparicio OM, Billington BL, Gottschling DE:

Modifiers of Position Effect Are Shared Between Telomeric and Silent Mating-Type Loci in S. cerevisiae.

Cell 1991, 66: 1279-1287. MEDLINE Cited by

Return to citation reference [1]

47. Braunstein M, Rose AB, Holmes SG, Allis CD, Broach JR:
Transcriptional Silencing in Yeast Is Associated with Reduced Nucleosome Acetylation. Genes Dev 1993, 7: 592-604. MEDLINE Cited by

Return to citation reference [1]

48. • Allshire RC, Javerzat J-P, Redhead NJ, Cranston G:

Position Effect Variegation at Fission Yeast Centromeres.

Cell 1994, 76: 157-169. MEDLINE Cited by

A convincing demonstration that $\overline{\text{PEV}}$ occurs in Schizosaccharomyces pombe , and that yeast centromeric regions can induce PEV. The central core of all three S. pombe centromeres was shown to induce reversible, but stable, transcriptional inactivation of inserted genes. Repression was correlated with unusual nucleosome spacing and a requirement that the core sequences be located at the centromere.

Return to citation reference [1] [2] [3]

49. Takahashi K, Murakami S, Chikashige Y, Funabiki H, Niwa O, Yanagida M:

A Low Copy Number Central Sequence with Strict Symmetry and Unusual Chromatin Structure in Fission Yeast Centromere.

Mol Biol Cell 1992, 3: 819-835. MEDLINE Cited by

Return to citation reference [1] [2]

50. • Rivier DH, Rine J:

An Origin of DNA Replication and a Transcription Silencer Require a Common Element. Science 1992, 256: 659-663. MEDLINE Cited by

A link between silencing and replication in yeast was strongly suggested by this study, in which silencing and replication were shown to require the same HMR sequence element. The authors propose a role for replication in the establishment of the silenced state, rather than its maintenance.

Return to citation reference [1]

51. • Micklem G, Rowley A, Harwood J, Nasmyth K, Diffley JFX:

Yeast Origin Recognition Complex is Involved in DNA Replication and Transcriptional Silencing.

Nature 1993, 366: 87-89. MEDLINE Cited by

See [52••].

Return to citation reference [1] [2] [3]

52. •• Foss M, McNally FJ, Laurenson P, Rine J:

Origin Recognition Complex (ORC) in Transcriptional Silencing and DNA Replication in S. cerevisiae.

Science 1993, 262: 1838-1844. MEDLINE Cited by

This paper and $[51 \bullet \bullet]$ show, by genetic analyses, that the ORC2 gene, which encodes a component of the origin recognition complex, plays an important role in HMR silencing. These studies forge a clear mechanistic link between replication and silencing, in conjunction with complementary biochemical evidence described in [53..]. Three models for ORC function in silencing are discussed in this reference; the evidence favors ORC recruitment of other components (e.g. SIRs) to silenced regions, rather than a strict requirement for replication initiation (also see [53••]).

Return to citation reference [1] [2] [3] [4] [5]

53. • Bell SP, Kobayashi R, Stillman B:

Yeast Origin Recognition Complex Functions in Transcription Silencing and DNA Replication.

Science 1993, 262: 1844-1849. MEDLINE Cited by

The authors present biochemical evidence for a link between replication and silencing. Also see [52••].

Return to citation reference [1] [2] [3] [4]

54. • Chien C-T, Buck S, Sternglanz R, Shore D:

Targeting of SIR1 Protein Establishes Transcriptional Silencing at HMLoci and Telomeres in Yeast.

Cell 1993, **75**: 531-541. MEDLINE Cited by

The sufficiency of SIR proteins in silencing was analyzed by 'tethering' GAL4-SIR fusions (via UAS elements) to sites that lacked normal binding elements for ORC, RAP and ABF. Only SIR1 was shown to be sufficient to establish or enhance silencing, and a deletion analysis suggested that ORC acts to recruit SIRs to silenced regions. Also see [52...].

Return to citation reference [1]

55. Rabl C:

Uber Zelltheilung.

Morphologisches Jahrbuch 1885, 10: 214-330.

Return to citation reference [1]

56. Hochstrasser M, Mathog D, Gruenbaum Y, Saumweber H, Sedat J:

Spatial Organisation of Chromosomes in the Salivary Gland Nuclei of Drosophila melanogaster.

J Cell Biol 1986, 102: 112-123. MEDLINE Cited by

Return to citation reference [1]

57. • Funabiki H, Hagan I, Uzawa S, Yanagida M:

Cell Cycle- Dependent Specific Positioning and Clustering of Centromeres and **Telomeres in Fission Yeast.**

J Cell Biol 1993, 121: 961-976. MEDLINE Cited by See [58•].

Return to citation reference [1] [2]

58. • Vourc'h C, Taruscio D, Boyle AL, Ward DC:
Cell Cycle- Dependent Distribution of Telomeres, Centromeres, and Chromosome-Specific Subsatellite Domains in the Interphase Nucleus of Mouse Lymphocytes.

Exp Cell Res 1993, 205: 142-151. ScienceDirect MEDLINE Cited by This paper and [57•] describe cytological analyses of the positioning of telomeres and centromeres. They show that these structures are clustered in interphase nuclei and that their nuclear distribution changes during the cell cycle. The functional significance of these observations needs to be determined by genetic and molecular analyses.

Return to citation reference [1] [2]

59. • Gilson E, Laroche T, Gasser SM:

Telomeres and the Functional Architecture of the Nucleus.

Trends Cell Biol 1993, 3: 128–134. <u>ScienceDirect Cited by</u>
A good review of the nuclear localization of telomeres in yeast, and of the proteins involved in telomere-telomere and telomere- nuclear membrane associations.

Return to citation reference [1] [2]

60. Wakimoto BT, Hearn MG:

The Effects of Chromosome Rearrangements on the Expression of Heterochromatic Genes in Chromosome 2L of Drosophila melanogaster.

Genetics 1990, 125: 141--154. MEDLINE Cited by

Return to citation reference [1] [2] [3] [4]

61. Eberl DF, Duyf BJ, Hilliker AJ:

The Role of Heterochromatin in the Expression of a Heterochromatic Gene, the *rolled*Locus of *Drosophila melanogaster*. *Genetics* 1993, **134**: 277– 292. MEDLINE Cited by

Return to citation reference [1] [2] [3]

62. • Wines DR, Henikoff S:

Somatic Instability of a DrosophilaChromosome.

Genetics 1992, 131: 683-691. MEDLINE Cited by

The first demonstration that PEV can affect inheritance functions; instability of a rearranged chromosome was shown to cause variegation of a white gene. Surprisingly, this phenotype behaved like the PEV of heterochromatic genes; additional heterochromatin enhanced the variegation in trans . Cytological analysis of the transmission defect suggests a role for heterochromatin and associated proteins in initiating and/or maintaining chromatid apposition.

Return to citation reference [1]

63. • Talbert PB, LeCiel DS, Henikoff S:

Modification of the DrosophilaHeterochromatic Mutation brown DrosophilaHeterochromatic Mutations.

Genetics 1994, 136: 559-571. MEDLINE Cited by

The importance of nuclear positioning, rather than strict heterochromatin/euchromatin differences, is suggested by the distribution of rearrangement breakpoints associated with suppression of the heterochromatin-induced PEV on the brown gene (see [60][61] for other examples). The 'looping' of heterochromatin into specific nuclear domains is invoked to explain these observations. The bw D variegation is also an example of 'trans -sensing' position effects (reviewed in [7]), which suggest a role in transcription for another aspect of nuclear organization, the association of genes on homologous chromosomes ('somatic pairing').

Return to citation reference [1] [2] [3]

64. Tower J, Karpen GH, Craig N, Spradling AC:

Preferential Transposition of *Drosophila*P Elements to Nearby Chromosomal Sites. Genetics 1993, 133: 347–359. MEDLINE Cited by

Return to citation reference [1]

65. Zhang P, Spradling AC:

Efficient and Dispersed Local P Element Transposition from *Drosophila*Females. *Genetics* 1993, **133**: 361–373. <u>MEDLINE Cited by</u>

Return to citation reference [1]

66. Hiraoka Y, Dernburg AF, Parmelee SJ, Rykowski MC, Agard DA, Sedat JW:

The Onset of Homologous

Chromosome Pairing During Drosophila melanogaster Embryogenesis.

J Cell Biol 1993, 120: 591- -600. MEDLINE Cited by

Return to citation reference [1]

67. Karpen GH, Schaeffer J, Laird CD:

A *Drosophila*rRNA Gene Located in Euchromatin Is Active in Transcription and Nucleolus Formation.

Genes Dev 1988, 2: 1745-1763. MEDLINE Cited by

Return to citation reference [1]

68. McKee BD, Karpen GH:

DrosophilaRibosomal RNA Genes Function As an X-Y Pairing Site During Male Meiosis. *Cell* 1990, **61**: 61-72. MEDLINE Cited by

Return to citation reference [1]

69. Gilson E, Roberge M, Giraldo R, Rhodes D, Gasser SM:

Distortion of the DNA Double Helix by RAP1 at Silencers and Multiple Telomeric Binding Sites.

J Mol Biol 1993, 231: 293-310. ScienceDirect MEDLINE Cited by

Return to citation reference [1]

70. Kyrion G, Boakye KA, Lustig AJ:

C-Terminal Truncation of RAP1 Results in the Deregulation of Telomere Size, Stability, and Function in Saccharomyces cerevisiae.

Mol Cell Biol 1992, 12: 5159-5173. MEDLINE Cited by

Return to citation reference [1]

71. • Kyrion G, Liu K, Liu C, Lustig AJ:

RAP1 and Telomere Structure Regulate Telomere Position Effects in Saccharomyces cerevisiae.

Genes Dev 1993, 7: 1146-1159. MEDLINE Cited by

This paper shows a functional role for RAP1 in yeast PEV; carboxy terminal deletions alleviated telomeric and HML silencing, and increased accessibility of this chromatin to a methylase.

Return to citation reference [1]

72. • Palladino F, Laroche T, Gilson E, Axelrod A, Pillus L, Gasser SM:
SIR3 and SIR4 Proteins Are Required for the Positioning and Integrity of Yeast
Telomeres.

Cell 1993, **75**: 543-555. MEDLINE Cited by

Monitoring the nuclear distribution of RAP1 protein in mutant cells suggests that both SIR3 and SIR4 are required for clustering of telomeres, as well as telomere localization to the membrane. Loss of SIR3/4 function resulted in telomere repeat shortening and chromosome instability.

Return to citation reference [1]

73. Gall JG, Cohen EH, Polan ML:

Repetitive Sequences in Drosophila.

Chromosoma 1971, 33: 319-344. MEDLINE Cited by

Return to citation reference [1]

74. Karpen GH, Spradling AC:

Reduced DNA Polytenization of a Minichromosome Region Undergoing Position-Effect Variegation in *Drosophila*.

Cell 1990, 63: 97-107. MEDLINE Cited by

Return to citation reference [1] [2] [3] [4] [5]

75. Miklos GL, Cotsell JN:

Chromosome Structure at Interfaces Between Major Chromatin Types: Alpha- and Beta- Heterochromatin.

Bioessays 1990, 12: 1-6. MEDLINE Cited by

Return to citation reference [1] [2]

76. Umbetova GH, Belyaeva ES, Baricheva EM, Zhimulev IF:

Cytogenetic and Molecular Aspects of Position Effect Variegation in *Drosophila melanogaster*. IV. Underreplication of Chromosomal Material As a Result of Gene Inactivation.

Chromosoma 1991, 101: 55-61. MEDLINE Cited by

Return to citation reference [1]

77. • Karpen GH, Spradling AC:

Analysis of

Subtelomeric Heterochromatin in the *Drosophila* Minichromosome *Dp1187* by Single P Element Insertional Mutagenesis.

Genetics 1992, 132: 737-753. MEDLINE Cited by

P-element mutagenesis of a minichromosome led to the identification a large domain of subtelomeric heterochromatin in Drosophila , which displays the repeated DNA, PEV induction, and under-representation (reduced DNA copy number in polytene nuclei), usually associated with centric heterochromatin. Diversity of PEV mechanisms is suggested by the observation that subtelomeric PEV is likely caused by transcriptional repression, while 220 kb away the PEV associated with the centric heterochromatin includes significant underrepresentation.

Return to citation reference [1] [2] [3] [4] [5] [6]

78. Yamamoto MT, Mitchelson A, Tudor M, O'Hare K, Davies JA, Miklos GL:

Molecular and Cytogenetic Analysis of the Heterochromatin-Euchromatin Junction Region of the *Drosophila melanogasterX* Chromosome Using Cloned DNA Sequences. *Genetics* 1990, **125**: 821–832. MEDLINE Cited by

Return to citation reference [1]

79. Levis RW:

Viable Deletions of a Telomere from a DrosophilaChromosome.

Cell 1989, **58**: 791-801. MEDLINE Cited by

Return to citation reference [1] [2]

80. • Ferguson BM, Fangman WL:

A Position Effect on the Time of Replication Origin Activation in Yeast.

Cell 1992, 68: 333-339. MEDLINE Cited by

This study conclusively demonstrates that late replication of a yeast origin is caused by positioning near the telomere. C 1-3 A telomeric repeats alone were not sufficient to induce late replication on a circular plasmid. Thus the position effect on replication timing depends on the chromatin structure or topology of chromosome ends, rather than telomere sequence composition.

Return to citation reference [1]

81. Laird CD:

DNA of DrosophilaChromosomes.

Annu Rev Genet 1973, 7: 177-204. MEDLINE Cited by

Return to citation reference [1]

82. • Hansen RS, Canfield TK, Lamb MM, Gartler SM, Laird CD:

Association of Fragile X Syndrome with Delayed Replication of the FMR1 Gene.

Cell 1993, 73: 1403-1409. MEDLINE Cited by

This paper presents clear evidence for altered replication timing of the FMR1 gene in human chromosomes at the Fragile X site. The authors describe a novel PCR technique for assessing replication in late versus early S phase.

Return to citation reference [1]

83. Pimpinelli S, Goday C:

Unusual Kinetochores and Chromatin Diminution in Parascaris.

Trends Genet 1989, 5: 310-315. ScienceDirect MEDLINE Cited by

Return to citation reference [1]

84. • Prescott DM:

Restructuring of DNA Sequences in the Germline Genome of Oxytricha.

Curr Opin Genet Dev 1993, 3: 726-729. MEDLINE Cited by

A concise review of the highly regulated DNA rearrangements and eliminations associated with macronuclear development in ciliates.

Return to citation reference [1] [2]

85. Spradling AC, Karpen G, Glaser R, Zhang P: **Evolutionary Conservation of Developmental Mechanisms: DNA Elimination in Drosophila.** In *phila* Edited by Spradling AC. New York: Wiley-Liss Inc, 1993, 39–53.

Return to citation reference [1] [2]

86. • Glaser RL, Karpen GH, Spradling AC:

Replication Forks Are Not Found in a *Drosophila* Minichromosome Demonstrating a Gradient of Polytenization.

Chromosoma 1992, 102: 15-19. MEDLINE Cited by

Two-dimensional gel electrophoresis was used to analyze intermediates present in a gradient of underrepresented DNA in a Drosophila minichromosome. Contrary to the predictions of the 'underreplication' model, stalled fork structures were not found, suggesting that heterochromatic underrepresentation in dipteran polytene cells may involve a DNA elimination mechanism.

Return to citation reference [1]

87. • Spradling AC:

Position Effect Variegation and Genomic Instability.

Cold Spring Harb Symp Quant Biol 1994, 58: in press. Cited by

Pulsed-field analysis demonstrates the occurrence of rearranged minichromosomes in Drosophila polytenized tissues. This represents positive evidence that heterochromatin elimination may occur in Drosophila . The author discusses potential roles for elimination in a variety of organisms.

Return to citation reference [1] [2]

88. Jaraczewski J, Jahn C:

Elimination of Tec Elements Involves a Novel Excision Process.

Genes Dev 1993, 7: 95-105. MEDLINE Cited by

Return to citation reference [1]

Author Contacts



GH Karpen, Molecular Biology and Virology Laboratory, The Salk Institute for Biological Studies, 10010 North Torrey Pines Road, La Jolla, California, 92037-1099, USA

Return to author list

Copyright

Copyright © 1994 Current Opinions



A Scientific Breakthrough

BioMedNetInformation for Advertisers

Journal Collection

Reviews

Research Update Research

News &

Books &

Science



© Elsevier Limited 2003